# Author Index


Abe, A., see Yamane, M. 579(1992)25

Abolfathi, Z., Bélanger, P.-M., Gilbert, M., Rouleau, J. R. and Turgeon, J.
Improved high-performance liquid chromatographic assay for the stereoselective determination of mexiletine in plasma 579(1992)366

Abramowski, V., see Bories, C. 579(1992)143

Akiyama, H., Shidawara, S., Mada, A., Toyoda, H., Toida, T. and Imanari, T.
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

Alton, K. B., Hernandez, A., Alvarez, N. and Patrick, J. E.
High-performance liquid chromatographic determination of N-[2(*S*)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

Alvarez, N., see Alton, K. B. 579(1992)307

Anderson, K. L., see Wang, L. H. 579(1992)259

Andreux, J. P., see Merino-Dugay, A. 579(1992)73

Aspeslet, L. J., see Torok-Both, G. A. 579(1992)99

Baker, G. B., see Torok-Both, G. A. 579(1992)99

Baron, B., see Bories, C. 579(1992)143

Beck, O., see Helander, A. 579(1992)340

Bélanger, P.-M., see Abolfathi, Z. 579(1992)366

Benet, L. Z., see Han, C. 579(1992)237

Bhatti, M. M. and Foster, R. T.
Stereospecific high-performance liquid chromatographic assay of metoprolol 579(1992)361

Bogusz, M.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

Borg, S., see Helander, A. 579(1992)340

Bories, C., Cardot, P. J. P., Abramowski, V., Poüs, C., Merino-Dugay, A., Baron, B. and Mougeot, G.
Elution mode of *Pneumocystis carinii* cysts in gravitational field-flow fractionation 579(1992)143

Borner, E., see Borner, K. 579(1992)285

Borner, K., Borner, E. and Lode, H.
Determination of sparfloxacin in serum and urine by high-performance liquid chromatography 579(1992)285

Bottiglieri, T., see Hyland, K. 579(1992)55

Branham, K. E., see Meng, Q. C. 579(1992)63

Breckenridge, W. C., see Kuksis, A. 579(1992)13

Brørs, O., see Holthe, M. R. 579(1992)376

Brunmark, P., Persson, P. and Skarping, G.
Determination of 4,4′-methylenedianiline in hydrolysed human urine by micro liquid chromatography with ultraviolet detection 579(1992)350

Cardot, P. J. P., see Bories, C. 579(1992)143

Cardot, P. J. P., see Merino-Dugay, A. 579(1992)73

Caron, J., see Trivier, J. M. 579(1992)269

Castagnoli, Jr., N., see Igarashi, K. 579(1992)277

Champney, T. H., Hanneman, W. H. and Nichols, M. A.
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Chari, G.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

Chiba, K., see Kobayashi, K. 579(1992)299

Constant, H., see Duthel, J. M. 579(1992)85

Coutts, R. T., see Torok-Both, G. A. 579(1992)99

Cragoe, Jr., E. J., see Holthe, M. R. 579(1992)376

Czok, M., see Merino-Dugay, A. 579(1992)73

Delucas, L. J., see Meng, Q. C. 579(1992)63

Demby, K. B., White, T. C. and Ghanayem, B. I.
Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection 579(1992)153

Desai, D. M. and Gal, J.
Reversed-phase high-performance liquid chromatographic separation of the stereoisomers of labetalol via derivatization with chiral and non-chiral isothiocyanate reagents 579(1992)165

Deutsch, J., Hegedus, L., Greig, N. H., Rapoport, S. I. and Soncrant, T. T.
Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography–mass spectrometry in rat plasma and brain 579(1992)93

Doležalová, M.
On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

Dunnett, M. and Harris, R. C.
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

Duthel, J. M., Constant, H., Vallon, J. J., Rochet, T. and Miachon, S.
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

Endoh, Y. S., Yoshimura, H., Sasaki, N., Ishihara, Y., Sasaki, H., Nakamura, S., Inoue, Y. and Nishikawa, M.
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

Eto, S., Noda, H. and Noda, A.
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β-cyclodextrin inclusion complexes 579(1992)253

# Author Index


Abe, A., see Yamane, M. 579(1992)25

Abolfathi, Z., Bélanger, P.-M., Gilbert, M., Rouleau, J. R. and Turgeon, J.
Improved high-performance liquid chromatographic assay for the stereoselective determination of mexiletine in plasma 579(1992)366

Abramowski, V., see Bories, C. 579(1992)143

Akiyama, H., Shidawara, S., Mada, A., Toyoda, H., Toida, T. and Imanari, T.
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

Alton, K. B., Hernandez, A., Alvarez, N. and Patrick, J. E.
High-performance liquid chromatographic determination of N-[2(*S*)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

Alvarez, N., see Alton, K. B. 579(1992)307

Anderson, K. L., see Wang, L. H. 579(1992)259

Andreux, J. P., see Merino-Dugay, A. 579(1992)73

Aspeslet, L. J., see Torok-Both, G. A. 579(1992)99

Baker, G. B., see Torok-Both, G. A. 579(1992)99

Baron, B., see Bories, C. 579(1992)143

Beck, O., see Helander, A. 579(1992)340

Bélanger, P.-M., see Abolfathi, Z. 579(1992)366

Benet, L. Z., see Han, C. 579(1992)237

Bhatti, M. M. and Foster, R. T.
Stereospecific high-performance liquid chromatographic assay of metoprolol 579(1992)361

Bogusz, M.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

Borg, S., see Helander, A. 579(1992)340

Bories, C., Cardot, P. J. P., Abramowski, V., Poüs, C., Merino-Dugay, A., Baron, B. and Mougeot, G.
Elution mode of *Pneumocystis carinii* cysts in gravitational field-flow fractionation 579(1992)143

Borner, E., see Borner, K. 579(1992)285

Borner, K., Borner, E. and Lode, H.
Determination of sparfloxacin in serum and urine by high-performance liquid chromatography 579(1992)285

Bottiglieri, T., see Hyland, K. 579(1992)55

Branham, K. E., see Meng, Q. C. 579(1992)63

Breckenridge, W. C., see Kuksis, A. 579(1992)13

Brørs, O., see Holthe, M. R. 579(1992)376

Brunmark, P., Persson, P. and Skarping, G.
Determination of 4,4′-methylenedianiline in hydrolysed human urine by micro liquid chromatography with ultraviolet detection 579(1992)350

Cardot, P. J. P., see Bories, C. 579(1992)143

Cardot, P. J. P., see Merino-Dugay, A. 579(1992)73

Caron, J., see Trivier, J. M. 579(1992)269

Castagnoli, Jr., N., see Igarashi, K. 579(1992)277

Champney, T. H., Hanneman, W. H. and Nichols, M. A.
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

Chari, G.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

Chiba, K., see Kobayashi, K. 579(1992)299

Constant, H., see Duthel, J. M. 579(1992)85

Coutts, R. T., see Torok-Both, G. A. 579(1992)99

Cragoe, Jr., E. J., see Holthe, M. R. 579(1992)376

Czok, M., see Merino-Dugay, A. 579(1992)73

Delucas, L. J., see Meng, Q. C. 579(1992)63

Demby, K. B., White, T. C. and Ghanayem, B. I.
Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection 579(1992)153

Desai, D. M. and Gal, J.
Reversed-phase high-performance liquid chromatographic separation of the stereoisomers of labetalol via derivatization with chiral and non-chiral isothiocyanate reagents 579(1992)165

Deutsch, J., Hegedus, L., Greig, N. H., Rapoport, S. I. and Soncrant, T. T.
Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography–mass spectrometry in rat plasma and brain 579(1992)93

Doležalová, M.
On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

Dunnett, M. and Harris, R. C.
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

Duthel, J. M., Constant, H., Vallon, J. J., Rochet, T. and Miachon, S.
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

Endoh, Y. S., Yoshimura, H., Sasaki, N., Ishihara, Y., Sasaki, H., Nakamura, S., Inoue, Y. and Nishikawa, M.
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

Eto, S., Noda, H. and Noda, A.
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β-cyclodextrin inclusion complexes 579(1992)253

Feather, T., see Kuksis, A. 579(1992)13
Fielding, R. M., see Wang, L. H. 579(1992)259
Foster, R. T., see Bhatti, M. M. 579(1992)361
Furuta, T., see Shibasaki, H. 579(1992)193
Gal, J., see Desai, D. M. 579(1992)165
Geher, K., see Kuksis, A. 579(1992)13
Ghanayem, B. I., see Demby, K. B. 579(1992)153
Gilbert, M., see Abolfathi, Z. 579(1992)366
Greig, N. H., see Deutsch, J. 579(1992)93
Guernet, M., see Merino-Dugay, A. 579(1992)73
Gumbleton, M., see Han, C. 579(1992)237
Hage, D. S., see Loun, B. 579(1992)225
Han, C., Gumbleton, M., Lau, D. T. W. and Benet, L. Z.
Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237
Hanneman, W. H., see Champney, T. H. 579(1992)334
Harris, R. C., see Dunnett, M. 579(1992)45
Hattori, H., Yamamoto, S., Iwata, M., Takashima, E., Yamada, T. and Suzuki, O.
Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247
Hegedus, L., see Deutsch, J. 579(1992)93
Helander, A., Beck, O. and Borg, S.
Determination of urinary 5-hydroxytryptophol by high-performance liquid chromatography with electrochemical detection 579(1992)340
Hernandez, A., see Alton, K. B. 579(1992)307
Holthe, M. R., Brørs, O., Cragoe, Jr., E. J., Veel, T. and Ræder, M.
Determination of active 5-(N,N-hexamethylene)amiloride in pig plasma 579(1992)376
Hyland, K. and Bottiglieri, T.
Measurement of total plasma and cerebrospinal fluid homocysteine by fluorescence following high-performance liquid chromatography and precolumn derivatization with *o*-phthaldialdehyde 579(1992)55
Ide, T., see Murata, M. 579(1992)329
Igarashi, K. and Castagnoli, Jr., N.
Determination of the pyridinium metabolite derived from haloperidol in brain tissue, plasma and urine by high-performance liquid chromatography with fluorescence detection 579(1992)277
Imanari, T., see Akiyama, H. 579(1992)203
Inoue, Y., see Endoh, Y. S. 579(1992)123
Ishihara, Y., see Endoh, Y. S. 579(1992)123
Ishikawa, F., see Yamane, M. 579(1992)25
Ishizaki, T., see Kobayashi, K. 579(1992)299
Iwata, M., see Hattori, H. 579(1992)247
Jansen, R. W., see Molema, G. 579(1992)107
Kasuya, Y., see Shibasaki, H. 579(1992)193
Kato, Y., see Kobayashi, K. 579(1992)299
Kema, I. P., see Muskiet, F. A. J. 579(1992)184
King, S. J., see Meng, Q. C. 579(1992)63
Kintz, P., Tracqui, A. and Mangin, P.
Determination of nalbuphine using high-performance liquid chromatography coupled to photodiode-array detection and gas chromatography coupled to mass spectrometry 579(1992)172
Knadler, M. P.
Quantification of a decapeptide anticoagulant in rat and monkey plasma by high-performance liquid chromatography 579(1992)371
Kobayashi, K., Chiba, K., Sohn, D.-R., Kato, Y. and Ishizaki, T.
Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated $C_{18}$ column 579(1992)299
Kodama, H., see Sugahara, K. 579(1992)318
Kolker, H. J., see Van 't Klooster, G. A. E. 579(1992)158
Kolker, H. J., see Van 't Klooster, G. A. E. 579(1992)355
Kuksis, A., Myher, J. J., Geher, K., Breckenridge, W. C., Feather, T., McGuire, V. and Little, J. A.
Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with carbohydrate and alcohol intake based on 24-h dietary recall 579(1992)13
Lau, D. T. W., see Han, C. 579(1992)237
Lhermitte, M., see Trivier, J. M. 579(1992)269
Li, K.
Determination of sialic acids in human serum by reversed-phase liquid chromatography with fluorimetric detection 579(1992)209
Libersa, C., see Trivier, J. M. 579(1992)269
Little, J. A., see Kuksis, A. 579(1992)13
Liu, H. and Worthen, H. G.
Measurement of free amino acid levels in ultrafiltrates of blood plasma by high-performance liquid chromatography with automatic pre-column derivatization 579(1992)215
Lode, H., see Borner, K. 579(1992)285
Lorber, B., see Meng, Q. C. 579(1992)63
Loun, B. and Hage, D. S.
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225
Mada, A., see Akiyama, H. 579(1992)203
Makino, Y. and Nishimura, S.
High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346
Mangin, P., see Kintz, P. 579(1992)172
Martín-Villacorta, J., see Méndez, R. 579(1992)115
McGuire, V., see Kuksis, A. 579(1992)13
McKenna, K. F., see Torok-Both, G. A. 579(1992)99
Meijer, D. K. F., see Molema, G. 579(1992)107
Méndez, R., Negro, A. and Martín-Villacorta, J.
High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115
Meng, Q. C., King, S. J., Branham, K. E., Delucas, L. J., Lorber, B. and Oparil, S.
Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63
Merino-Dugay, A., see Bories, C. 579(1992)143
Merino-Dugay, A., Cardot, P. J. P., Czok, M., Guernet, M. and Andreux, J. P.
Monitoring of an experimental red blood cell pathology with gravitational field-flow fractionation 579(1992)73

Miachon, S., see Duthel, J. M. 579(1992)85
Molema, G., Jansen, R. W., Visser, J. and Meijer, D. K. F.
Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107
Mougeot, G., see Bories, C. 579(1992)143
Muh, J. P.
Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites. Reply to Muskiet and Kema 579(1992)187
Murata, M. and Ide, T.
Determination of cholesterol in sub-nanomolar quantities in biological fluids by high-performance liquid chromatography 579(1992)329
Muskiet, F. A. J. and Kema, I. P.
Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites 579(1992)184
Myher, J. J., see Kuksis, A. 579(1992)13
Nakamura, S., see Endoh, Y. S. 579(1992)123
Negro, A., see Méndez, R. 579(1992)115
Nichols, M. A., see Champney, T. H. 579(1992)334
Nishikawa, M., see Endoh, Y. S. 579(1992)123
Nishimura, S., see Makino, Y. 579(1992)346
Noda, A., see Eto, S. 579(1992)253
Noda, H., see Eto, S. 579(1992)253
Noordhoek, J., see Van 't Klooster, G. A. E. 579(1992)158
Noordhoek, J., see Van 't Klooster, G. A. E. 579(1992)355
Nouws, J. F. M., see Vree, T. B. 579(1992)131
Ohta, J., see Sugahara, K. 579(1992)318
Oparil, S., see Meng, Q. C. 579(1992)63
Patrick, J. E., see Alton, K. B. 579(1992)307
Pellegrino, S., see Petrarulo, M. 579(1992)324
Persson, P., see Brunmark, P. 579(1992)350
Petrarulo, M., Pellegrino, S. and Testa, E.
High-performance liquid chromatographic microassay for methyl ethyl ketone in urine as the 2,4-dinitrophenylhydrazone derivative 579(1992)324
Pommery, J., see Trivier, J. M. 579(1992)269
Poüs, C., see Bories, C. 579(1992)143
Ræder, M., see Holthe, M. R. 579(1992)376
Rapoport, S. I., see Deutsch, J. 579(1992)93
Rochet, T., see Duthel, J. M. 579(1992)85
Rouleau, J. R., see Abolfathi, Z. 579(1992)366
Rubino, F. M. and Zecca, L.
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1
Sasaki, H., see Endoh, Y. S. 579(1992)123
Sasaki, N., see Endoh, Y. S. 579(1992)123
Shearer, G., see Tarbin, J. A. 579(1992)177
Shibasaki, H., Furuta, T. and Kasuya, Y.
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193
Shidawara, S., see Akiyama, H. 579(1992)203
Skarping, G., see Brunmark, P. 579(1992)350
Smith, P. C., see Wang, L. H. 579(1992)259
Sohn, D.-R., see Kobayashi, K. 579(1992)299
Soncrant, T. T., see Deutsch, J. 579(1992)93
Spaethe, S. M., see VanAlstyne, E. L. 579(1992)37
Sugahara, K., Ohta, J., Takemura, M. and Kodama, H.
Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography–mass spectrometry 579(1992)318
Suzuki, O., see Hattori, H. 579(1992)247
Takashima, E., see Hattori, H. 579(1992)247
Takemura, M., see Sugahara, K. 579(1992)318
Tarbin, J. A. and Shearer, G.
Improved high-performance liquid chromatographic procedure for the determination of lasalocid in chicken tissues and egg using polymeric and porous graphitic carbon columns 579(1992)177
Testa, E., see Petrarulo, M. 579(1992)324
Toida, T., see Akiyama, H. 579(1992)203
Torok-Both, G. A., Baker, G. B., Coutts, R. T., McKenna, K. F. and Aspeslet, L. J.
Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99
Toyoda, H., see Akiyama, H. 579(1992)203
Tracqui, A., see Kintz, P. 579(1992)172
Trivier, J. M., Pommery, J., Libersa, C., Caron, J. and Lhermitte, M.
High-performance liquid chromatographic assay for amiodarone N-deethylation in microsomes of rat liver 579(1992)269
Turgeon, J., see Abolfathi, Z. 579(1992)366
Vallon, J. J., see Duthel, J. M. 579(1992)85
VanAlstyne, E. L. and Spaethe, S. M.
Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37
Van Ewijk-Beneken Kolmer, E. W. J., see Vree, T. B. 579(1992)131
Van Miert, A. S. J. P. A. M., see Van 't Klooster, G. A. E. 579(1992)158
Van Miert, A. S. J. P. A. M., see Van 't Klooster, G. A. E. 579(1992)355
Van 't Klooster, G. A. E., Kolker, H. J., Woutersen-van Nijnanten, F. M. A., Noordhoek, J. and Van Miert, A. S. J. P. A. M.
Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355
Van 't Klooster, G. A. E., Woutersen-van Nijnanten, F. M. A., Kolker, H. J., Noordhoek, J. and Van Miert, A. S. J. P. A. M.
Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158
Veel, T., see Holthe, M. R. 579(1992)376
Visser, J., see Molema, G. 579(1992)107

Vree, T. B., Van Ewijk-Beneken Kolmer, E. W. J. and Nouws, J. F. M.
Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131

Wang, L. H., Smith, P. C., Anderson, K. L. and Fielding, R. M.
High-performance liquid chromatographic analysis of amphotericin B in plasma, blood, urine and tissues for pharmacokinetic and tissue distribution studies 579(1992)259

White, T. C., see Demby, K. B. 579(1992)153

Worthen, H. G., see Liu, H. 579(1992)215

Woutersen-van Nijnanten, F. M. A., see Van 't Klooster, G. A. E. 579(1992)158

Woutersen-van Nijnanten, F. M. A., see Van 't Klooster, G. A. E. 579(1992)355

Yamada, T., see Hattori, H. 579(1992)247

Yamamoto, S., see Hattori, H. 579(1992)247

Yamane, M., Abe, A., Yamane, S. and Ishikawa, F.
High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

Yamane, S., see Yamane, M. 579(1992)25

Yoshimura, H., see Endoh, Y. S. 579(1992)123

Zecca, L., see Rubino, F. M. 579(1992)1

# Compound Index

**N-Acetyldopamine**
Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites 579(1992)184

Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites. Reply to Muskiet and Kema 579(1992)187

**Acids**
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

**Albumin**
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

**Alcohols**
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

**Amino acids**
Measurement of free amino acid levels in ultrafiltrates of blood plasma by high-performance liquid chromatography with automatic pre-column derivatization 579(1992)215

**γ-Aminobutyric acid**
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Amiodarone**
High-performance liquid chromatographic assay for amiodarone N-deethylation in microsomes of rat liver 579(1992)269

**Amitriptyline**
On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

**Amphotericin B**
High-performance liquid chromatographic analysis of amphotericin B in plasma, blood, urine and tissues for pharmacokinetic and tissue distribution studies 579(1992)259

**Angiotensin-converting enzyme**
Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63

**Anserine**
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

**Apolactoferrin**
High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346

**Atriopeptidase**
High-performance liquid chromatographic determination of N-[2(*S*)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

**Azidothymidine**
Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107

**Azidothymidine phosphates**
Simultaneous analysis of azidothymidine and its mono-, di- and triphosphate derivatives in biological fluids, tissue and cultured cells by a rapid high-performance liquid chromatographic method 579(1992)107

**Barbiturates**
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via β-cyclodextrin inclusion complexes 579(1992)253

**Carboxyprimaquine**
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

**Carnosine**
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

**Catecholamines**
Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites 579(1992)184

Sensitivity and specificity of urinary N-acetyldopamine as a marker for neuroblastomas: comparison with traditional urinary catecholamine metabolites. Reply to Muskiet and Kema 579(1992)187

γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Cholesterol**
Determination of cholesterol in sub-nanomolar quantities in biological fluids by high-performance liquid chromatography 579(1992)329

**Chondroitin sulphate**
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

**Codeine**
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

**Corticosteroids**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Cortisol**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Cortisone**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Cotinine**
Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography–mass spectrometry in rat plasma and brain 579(1992)93

**Cystathionine**
Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography–mass spectrometry 579(1992)318

**Decapeptides**
Quantification of a decapeptide anticoagulant in rat and monkey plasma by high-performance liquid chromatography 579(1992)371

**Dermatan sulphate**
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

**N-Desethylamiodarone**
High-performance liquid chromatographic assay for amiodarone N-deethylation in microsomes of ţat liver 579(1992)269

**N-Desmethyldiazepam**
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

**Diazepam**
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

**Dopamine**
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Enzymes**
Preparative isolation of angiotensin-converting enzyme from human lung 579(1992)63

**Ethylmorphine**
Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

**Fatty acids**
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

**Fatty alcohols**
Application of triple quadrupole tandem mass spectrometry to the analysis of pyridine-containing derivatives of long-chain acids and alcohols 579(1992)1

**Flumequine**
Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131

**Flumequine acyl glucuronide**
Direct-gradient high-performance liquid chromatographic analysis and preliminary pharmacokinetics of flumequine and flumequine acyl glucuronide in humans: effect of probenecid 579(1992)131

**Fluoxetine**
Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99

**Glyceryl dinitrate**
Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

**Glyceryl mononitrate**
Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

**Glycosaminoglycans**
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

**Haloperidol pyridinium metabolite**
Determination of the pyridinium metabolite derived from haloperidol in brain tissue, plasma and urine by high-performance liquid chromatography with fluorescence detection 579(1992)277

**5-(N,N-Hexamethylene)amiloride**
Determination of active 5-(N,N-hexamethylene)amiloride in pig plasma 579(1992)376

**Histidine**
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

**Homocysteine**
Measurement of total plasma and cerebrospinal fluid homocysteine by fluorescence following high-performance liquid chromatography and precolumn derivatization with *o*-phthaldialdehyde 579(1992)55

**Human serum albumin**
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

**Hyaluronic acid**
Chemiluminescence high-performance liquid chromatography for the determination of hyaluronic acid, chondroitin sulphate and dermatan sulphate 579(1992)203

**Hydantoin**
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via *β*-cyclodextrin inclusion complexes 579(1992)253

**Hydroperoxy polyunsaturated fatty acids**
High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

**12-Hydroxy-5,8,10,14-eicosatetraenoic acid**
Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37

**5*R*,6*S*,8*R*-6-(1-Hydroxyethyl)-2-carbamoyloxymethylpenem-3-carboxylic acid**
High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

**5*R*,6*R*,8*R*-6-(1-Hydroxyethyl)-2-ethylthiopenem-3-carboxylic acid**
High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

**5-Hydroxyindoleacetic acid**
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Hydroxyomeprazole**
Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated $C_{18}$ column 579(1992)299

**5-(*p*-Hydroxyphenyl)-5-phenylhydantoin**
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via *β*-cyclodextrin inclusion complexes 579(1992)253

**Hydroxytrimethoprim**
Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

**5-Hydroxytryptophol**
Determination of urinary 5-hydroxytryptophol by high-performance liquid chromatography with electrochemical detection 579(1992)340

**Imidazoles**
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

**Indole**
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

**Labetalol**
Reversed-phase high-performance liquid chromatographic separation of the stereoisomers of labetalol via derivatization with chiral and non-chiral isothiocyanate reagents 579(1992)165

**Lactoferrins**
High-performance liquid chromatographic separation of human apolactoferrin and monoferric and diferric lactoferrins 579(1992)346

**Lasalocid**
Improved high-performance liquid chromatographic procedure for the determination of lasalocid in chicken tissues and egg using polymeric and porous graphitic carbon columns 579(1992)177

**Leukotrienes**
Rapid method for automated on-line extraction and fractionation of plasma leukotrienes and 12-hydroxy-5,8,10,14-eicosatetraenoic acids by reversed-phase high-performance liquid chromatography 579(1992)37

**Lipids**
Gas chromatographic profiles of plasma total lipids as indicators of dietary history. Correlation with carbohydrate and alcohol intake based on 24-h dietary recall 579(1992)13

Determination of cholesterol in sub-nanomolar quantities in biological fluids by high-performance liquid chromatography 579(1992)329

**N-[2(*S*)-(Mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine**
High-performance liquid chromatographic determination of N-[2(*S*)-(mercaptomethyl)-3-(2-methylphenyl)-1-oxopropyl]-L-methionine, the active plasma metabolite of a prodrug atriopeptidase inhibitor (SCH 42495), using a thiol selective (Au/Hg) amperometric detector 579(1992)307

**Methacrylonitrile**
Determination of methacrylonitrile in serum using gas chromatography and flame ionization detection 579(1992)153

**4,4′-Methylenedianiline**
Determination of 4,4′-methylenedianiline in hydrolysed human urine by micro liquid chromatography with ultraviolet detection 579(1992)350

**Methyl ethyl ketone**
High-performance liquid chromatographic microassay for methyl ethyl ketone in urine as the 2,4-dinitrophenylhydrazone derivative 579(1992)324

**1-Methylhistidine**
Determination of carnosine and other biogenic imidazoles in equine plasma by isocratic reversed-phase ion-pair high-performance liquid chromatography 579(1992)45

**Methylphenobarbital**
Chiral separation of barbiturates and hydantoins by reversed-phase high-performance liquid chromatography using a 25 or 50 mm short ODS cartridge column via $\beta$-cyclodextrin inclusion complexes 579(1992)253

**Metoprolol**
Stereospecific high-performance liquid chromatographic assay of metoprolol 579(1992)361

**Mexiletine**
Improved high-performance liquid chromatographic assay for the stereoselective determination of mexiletine in plasma 579(1992)366

**Morphine**
Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

**Morphine glucuronides**
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 579(1992)189

High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Bogusz 579(1992)191

**Nalbuphine**
Determination of nalbuphine using high-performance liquid chromatography coupled to photodiode-array detection and gas chromatography coupled to mass spectrometry 579(1992)172

**Nicotine**
Electron-impact and chemical ionization detection of nicotine and cotinine by gas chromatography–mass spectrometry in rat plasma and brain 579(1992)93

**Nitroglycerin**
Improved gas chromatographic assay for the simultaneous determination of nitroglycerin and its mono- and dinitrate metabolites 579(1992)237

**Norepinephrine**
$\gamma$-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Norethylmorphine**
Improved high-performance liquid chromatographic method for the determination of ethylmorphine and its metabolites in microsomal incubations and cell culture media 579(1992)158

**Norfluoxetine**
Simultaneous determination of fluoxetine and norfluoxetine enantiomers in biological samples by gas chromatography with electron-capture detection 579(1992)99

**Nortriptyline**
On-line solid-phase extraction and high-performance liquid chromatographic determination of nortriptyline and amitriptyline in serum 579(1992)291

**Omeprazole**
Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated $C_{18}$ column 579(1992)299

**Omeprazole sulphone**
Simultaneous determination of omeprazole and its metabolites in plasma and urine by reversed-phase high-performance liquid chromatography with an alkaline-resistant polymer-coated $C_{18}$ column 579(1992)299

**Oxazepam**
Quantitation by gas chromatography with selected-ion monitoring mass spectrometry of "natural" diazepam, N-desmethyldiazepam and oxazepam in normal human serum 579(1992)85

**Pamaquine**
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

**Penems**
High-performance liquid chromatographic methods for the determination of the penems SCH 29482 and FCE 22101 in human serum and urine 579(1992)115

**Perhydro-1,4-thiazepine-3,5-dicarboxylic acid**
Determination of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using column liquid chromatography–mass spectrometry 579(1992)318

**Phenothiazines**
Sensitive determination of phenothiazines in body fluids by gas chromatography with surface ionization detection 579(1992)247

**Polyunsaturated fatty acids**
High-performance liquid chromatography–thermospray mass spectrometry of hydroperoxy polyunsaturated fatty acid acetyl derivatives 579(1992)25

**Prednisolone**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Prednisone**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Primaquine**
High-performance liquid chromatographic determination of pamaquine, primaquine and carboxy primaquine in calf plasma using electrochemical detection 579(1992)123

**Serotonin**
γ-Aminobutyric acid, catecholamine and indoleamine determinations from the same brain region by high-performance liquid chromatography with electrochemical detection 579(1992)334

**Sialic acids**
Determination of sialic acids in human serum by reversed-phase liquid chromatography with fluorimetric detection 579(1992)209

**Sparfloxacin**
Determination of sparfloxacin in serum and urine by high-performance liquid chromatography 579(1992)285

**Steroids**
Stable isotope dilution mass spectrometry for the simultaneous determination of cortisol, cortisone, prednisolone and prednisone in plasma 579(1992)193

**Thyroxine**
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225

**Trimethoprim**
Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

**Trimethoprim N-oxides**
Determination of trimethoprim and its oxidative metabolites in cell culture media and microsomal incubation mixtures by high-performance liquid chromatography 579(1992)355

**Warfarin**
Characterization of thyroxine–albumin binding using high-performance affinity chromatography. I. Interactions at the warfarin and indole sites of albumin 579(1992)225